



CrossMark

In this issue

doi:10.1136/acupmed-2017-011556

David John Carr^{1,2}

This issue of *Acupuncture in Medicine* opens with two new systematic reviews with a decidedly gastrointestinal theme that explore the potentially positive effects of acupuncture at opposite ends of the alimentary tract. The first, by Zhu *et al* pooled the results of 12 randomised controlled trials of manual acupuncture (MA) and electroacupuncture (EA) as both adjuncts and alternatives to conventional medications (predominantly proton pump inhibitors) to treat gastro-oesophageal reflux disease (GORD). Compared with pharmacological therapy alone, recurrence rates of GORD were significantly lower in MA/EA-treated patients and, when MA/EA was combined with medication, there was a greater global symptom improvement. They also found evidence of a beneficial effect on quality of life measures, suggesting this is an area worthy of further investigation, especially given the relatively limited methodological quality of the existing evidence. The second systematic review, by Zhou *et al*, presents a comparison of the effects of EA versus conventional medications (eg, mosapride, senna and lactulose) for functional constipation. EA treatment was associated with a greater frequency of spontaneous bowel movements, larger reductions in symptom scores and higher total response rates. Their findings are consistent with recent preclinical research in a cold saline gavage rat model of constipation, in which objective improvements in gastrointestinal transit and defaecation frequency were accompanied by enhanced colonic serotonin and tryptophan hydroxy-lase expression.¹

Next, Zhao *et al* present the findings of a multicentre randomised controlled trial of acupuncture at SP6 for primary dysmenorrhoea. This study was originally designed to test the importance of the *de qi* sensation in treatment outcome, which is pertinent given that it has been recently shown that many determinants of acupuncture 'dose', including the number of needles used, retention time and overall number of treatments, have a significant impact on therapeutic outcomes in

this particular condition.² It therefore seems feasible that the degree of manual needle stimulation (ie, elicitation of *de qi*) may also be important in this respect. The authors found however, that *prevention* of *de qi* sensation was much harder than expected, given that 55% of participants who were randomised to the 'no *de qi*' group still experienced the classical sensation. Accordingly, intention-to-treat analysis resulted in no significant difference between groups. Interestingly, however, actual experience of *de qi*, independent of original group allocation was associated with apparently greater analgesic effects.

It is increasingly recognised that acupuncture and related techniques impact the autonomic nervous system.^{3,4} Waki *et al* add further to this literature by showing that high-frequency (100Hz) facial EA increases parasympathetic activity as well as enhances cerebral blood flow. Elsewhere, Weeks *et al* highlight the potential receptivity of patients to the prospect of receiving acupuncture to prevent post-operative nausea and vomiting, the efficacy and effectiveness of which is comparable to anti-emetics,⁵ and which continues to be a focus of recent research evaluating the role of acupuncture in multi-modal post-surgical care.⁶⁻⁹

The issue concludes with four preclinical studies examining the effects of MA/EA on glucose levels and markers of ovarian function in female rats subjected to continuous light exposure, on hypoxia-inducible factor 1 α expression and infarct volume in male rats exposed to middle cerebral artery occlusion-induced ischaemic stroke, on the endoplasmic reticulum stress response in male rats with heroin addiction/relapse, and on urinary metabolomics markers in spontaneously hypertensive rats. The heroin extends the author's previous work in this area that evaluated the impact of acupuncture on apoptosis and the unfolded protein response.¹⁰

Competing interests None declared.

Provenance and peer review Not commissioned; internally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Zhu X, Liu Z, Qu H, *et al*. The effect and mechanism of electroacupuncture at LI11 and ST37 on constipation in a rat model. *Acupunct Med* 2016;34:194–200.

- Armour M, Smith CA. Treating primary dysmenorrhoea with acupuncture: a narrative review of the relationship between acupuncture 'dose' and menstrual pain outcomes. *Acupunct Med* 2016;34:416–24.
- Hideaki W, Tatsuya H, Shogo M, *et al*. Effect of 100 Hz electroacupuncture on salivary immunoglobulin A and the autonomic nervous system. *Acupunct Med* 2015;33:451–6.
- de la Cruz Torres B, Albornoz Cabello M, García Bermejo P, *et al*. Autonomic responses to ultrasound-guided percutaneous needle electrolysis of the patellar tendon in healthy male footballers. *Acupunct Med* 2016;34:275–9.
- Lee A, Chan SK, Fan LT. Stimulation of the wrist acupuncture point PC6 for preventing postoperative nausea and vomiting. *Cochrane Database Syst Rev* 2015;2:CD003281.
- Kim KH, Kim DH, Kim HY, *et al*. Acupuncture for recovery after surgery in patients undergoing colorectal cancer resection: a systematic review and meta-analysis. *Acupunct Med* 2016;34:248–56.
- Liudden I, Sandvik L, Valeberg BT, *et al*. Acupuncture versus usual care for postoperative nausea and vomiting in children after tonsillectomy/adenoidectomy: a pragmatic, multicentre, double-blinded, randomised trial. *Acupunct Med* 2015;33:196–203.
- Dalamagka M, Mavrommatis C, Grosomanidis V, *et al*. Postoperative analgesia after low-frequency electroacupuncture as adjunctive treatment in inguinal hernia surgery with abdominal wall mesh reconstruction. *Acupunct Med* 2015;33:360–7.
- Tseng CC, Tseng A. Effect of acupuncture on postoperative adhesive intestinal obstruction. *Acupunct Med* 2015;33:338–9.
- Zhang Y, Cai XH, Zhang RJ, *et al*. Acupuncture regulates the unfolded protein response and inhibits apoptosis in a rat model of heroin relapse. *Acupunct Med* 2016;34:441–8.

¹Department of Obstetrics, Gynecology and Reproductive Sciences, The Robert Larner, M.D. College of Medicine, The University of Vermont, Burlington, Vermont, USA

²Department of Maternal and Fetal Medicine, UCL EGA Institute for Women's Health, University College London, London, UK

Correspondence to Dr David John Carr, Department of Obstetrics, Gynecology and Reproductive Sciences, University of Vermont Medical Center, 111 Colchester Avenue, Burlington, VT 05401, USA; dcarr@uvm.edu



In this issue

David John Carr

Acupunct Med 2017 35: 315
doi: 10.1136/acupmed-2017-011556

Updated information and services can be found at:
<http://aim.bmj.com/content/35/5/315>

These include:

References

This article cites 10 articles, 9 of which you can access for free at:
<http://aim.bmj.com/content/35/5/315#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://www.bmj.com/company/products-services/rights-and-licensing/>

To order reprints go to:
<http://journals.bmj.com/content/subscribers>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>