

Implementation of shared medical appointments to offer battlefield acupuncture efficiently to veterans with pain

BACKGROUND

The use of prescription opioids in the treatment of pain has increased notably over recent decades.¹ With this increase, dramatic unintended consequences have arisen. Rates of death from prescribed opioids increased fourfold between 2000 and 2014.² Integrative care has been suggested as a potentially safer alternative to opioids in the treatment of chronic non-cancer pain and acupuncture has been shown to be an effective treatment for chronic pain.³ Battlefield acupuncture (BFA), an easily learnt subset of auricular acupuncture, has been proposed to treat a variety of painful disorders in active military members and veterans.⁴ Patients undergo insertion of five auricular semi-permanent (ASP) needles to the following traditional ear acupuncture points bilaterally: *Cingulate Gyrus*, *Thalamus*, *Omega 2*, *Point Zero* and *Shenmen* (figure 1). While other investigators have evaluated models for integrating medical acupuncture into practice,⁵ to our knowledge, there has been no evaluation of how to incorporate BFA efficiently into a busy primary care (PC) practice. Since shared medical appointments (SMA) have been shown to be helpful in chronic disease management and may decrease health-care utilisation, we developed an SMA approach to deliver BFA in the setting of a US Department of Veterans Affairs (VA) PC practice.

IMPLEMENTING SMA FOR BFA IN PC

One of the authors (DGF) underwent 4 hours of BFA training by a certified medical acupuncturist (also trained in BFA) and, in an



Figure 1 Image of patient receiving battlefield acupuncture at the following traditional ear acupuncture points bilaterally: *Cingulate Gyrus*, *Thalamus*, *Omega 2*, *Point Zero* and *Shenmen*.

effort to expand the availability of the service and minimise any impact on access to routine care, we created a 90 min SMA solely for BFA. Patients were recruited via a large informational poster in the PC clinic waiting area that provided an explanation of BFA and invited them to ask their provider or nurse for more information, if interested. Simultaneously, emails were sent to all 36 PC providers at the West Haven VA medical centre informing them that an SMA was available to any of their patients who were interested in BFA for any acute or chronic pain. A separate 90 min group clinic was created within the VA's software package, capable of accommodating eight patients. Over the ensuing 4 weeks, seven additional clinics were created over a 3-month period and all 52 appointment slots were filled. To date, six group clinics have been completed and 43 patients have been seen within that context.

The first 30 min of the appointment consisted of a group informational session detailing what BFA entails, including potential risks and benefits, and aftercare.

It was also explained to patients that their pain issues needed to have been thoroughly assessed by their PC providers before they could receive BFA. Those willing to undergo the procedure then signed informed consent. During each clinic, the patient's baseline pre-treatment numerical pain score was recorded using the Defense and Veterans Pain Rating Scale (where 1='hardly notice pain' and 10='as bad as it could be, nothing else matters'). The pain score was recorded after each successive BFA point. Patients were contacted by telephone 1 day and 7 days after the SMA to assess their pain at that point in time.

FINDINGS

As shown in figure 2, the mean±SD pain scores at the start were 7.3±2.0. After BFA treatment, the pain scores decreased to 4.5±2.3 ($p<0.01$ using paired two sample t-test). At day 1 and day 7, the mean pain scores were 4.4±2.4 and 5.7±2.6, respectively ($p<0.01$ for both comparisons when compared with baseline). Pain decreased in 41 of 43 patients (95.3%) on the day of




Figure 2 Pain scores before, immediately after, and 1 and 7 days following a session of battlefield acupuncture (BFA) in 43 patients seen via a shared medical appointment model in a US Department of Veterans Affairs primary care practice. Data are presented as mean \pm SD.

BFA treatment, and in 39 of 43 patients (90.7%) on day 1. Pain scores remained lower in 30 of 42 patients (71.4%) on day 7. No patients reported increased pain with BFA on the day of treatment or 1 day after treatment. Only six of 42 patients contacted on day 7 (14.3%) reported increased pain 7 days after BFA treatment (

IMPLICATIONS FOR CLINICAL EFFECTIVENESS

Pain is associated with decreased quality of life and is one of the most common presenting symptoms in PC. However, it remains poorly understood and traditional, pharmacologically-based treatment approaches are fraught with complications. Thus, offering patients effective non-pharmacological options for pain relief should be a priority. BFA has the potential to help patients but, given the high prevalence of patients with chronic non-cancer pain, competing

medical comorbidities and the need to maintain access to healthcare providers, implementation of novel, non-traditional venues should be considered. We have described a well-received and efficient manner in which to offer an integrative modality within the context of a busy PC practice. Potential barriers that we were able to overcome included finding adequate space for a group of this size, obtaining training in BFA and institutional credentialing, and developing an efficient manner in which to perform BFA. Ultimately, our clinical results are limited by our self-selected patient population and the lack of a comparison group and longer term follow-up to ascertain the durability of response. However, our preliminary results suggest that BFA performed in a group setting at a VA institution appears effective for short-term pain reduction.

Daniel Glenn Federman,^{1,2} Lorie M Poulin,¹ Christopher B Ruser,^{1,2} Jeffrey D Kravetz^{1,2}

¹Department of Medicine, VA Connecticut Health System West Haven Campus, West Haven, Connecticut, USA

²Department of Medicine, Yale University School of Medicine, New Haven, Connecticut, USA

Correspondence to Dr Daniel Glenn Federman, Department of medicine, VA Connecticut Health System West Haven Campus, VA Connecticut 11ACSL 950 Campbell Ave. West Haven, CT 06516 ; daniel.federman@va.gov

Contributors All authors have contributed sufficiently in the design of our work and the manuscript preparation to warrant authorship.

Competing interests None declared.

Patient consent We were exempted by our IRB.

Ethics approval VA Connecticut IRB.

Provenance and peer review Not commissioned; internally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.



CrossMark

To cite Federman DG, Poulin LM, Ruser CB, et al. *Acupunct Med* Published Online First: [please include Day Month Year]. doi:10.1136/acupmed-2016-011315

Accepted 29 May 2017

Acupunct Med 2017;**0**:1–2.

doi:10.1136/acupmed-2016-011315

REFERENCES

- 1 Alford DP. Opioid prescribing for chronic pain--achieving the right balance through education. *N Engl J Med* 2016;374:301–3.
- 2 Compton WM, Jones CM, Baldwin GT. Relationship between nonmedical prescription-opioid use and heroin use. *N Engl J Med* 2016;374:154–63.
- 3 Vickers AJ, Cronin AM, Maschino AC, et al. Acupuncture for chronic pain: individual patient data meta-analysis. *Arch Intern Med* 2012;172:1444–53.
- 4 King HC, Hickey AH, Connelly C. Auricular acupuncture: a brief introduction for military providers. *Mil Med* 2013;178:867–74.
- 5 Crumley ET. Models for integrating medical acupuncture into practice: an exploratory qualitative study of physicians' experiences. *Acupunct Med* 2016;34:280–9.



Implementation of shared medical appointments to offer battlefield acupuncture efficiently to veterans with pain

Daniel Glenn Federman, Lorie M Poulin, Christopher B Ruser and Jeffrey D Kravetz

Acupunct Med published online June 19, 2017

Updated information and services can be found at:

<http://aim.bmj.com/content/early/2017/06/19/acupmed-2016-011315>

These include:

References

This article cites 5 articles, 1 of which you can access for free at:

<http://aim.bmj.com/content/early/2017/06/19/acupmed-2016-011315#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://www.bmj.com/company/products-services/rights-and-licensing/>

To order reprints go to:

<http://journals.bmj.com/content/subscribers>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>